

OFFICE OF NAVAL RESEARCH

Contract N00014-86-K-0043

TECHNICAL REPORT No. 119

Squeezing of Many-Atom Radiation in an Optical Cavity

by

Fu-li Li, Xiao-shen Li, D. L. Lin and Thomas F. George

Prepared for Publication

in

Physical Review A

Departments of Chemistry and Physics
State University of New York at Buffalo
Buffalo, New York 14260

November 1989

Reproduction in whole or in part is permitted for any purpose of the
United States Government.

This document has been approved for public release and sale;
its distribution is unlimited.

DTIC
ELECTE
DEC 12 1989

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

1a. REPORT SECURITY CLASSIFICATION Unclassified			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution unlimited		
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE					
4. PERFORMING ORGANIZATION REPORT NUMBER(S) UBUFFALO/DC/89/TR-119			5. MONITORING ORGANIZATION REPORT NUMBER(S)		
6a. NAME OF PERFORMING ORGANIZATION Depts. Chemistry & Physics State University of New York		6b. OFFICE SYMBOL (if applicable)		7a. NAME OF MONITORING ORGANIZATION	
6c. ADDRESS (City, State, and ZIP Code) Fronczak Hall, Amherst Campus Buffalo, New York 14260			7b. ADDRESS (City, State, and ZIP Code) Chemistry Program 800 N. Quincy Street Arlington, Virginia 22217		
8a. NAME OF FUNDING/SPONSORING ORGANIZATION Office of Naval Research		8b. OFFICE SYMBOL (if applicable)		9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER Contract N00014-86-K-0043	
8c. ADDRESS (City, State, and ZIP Code) Chemistry Program 800 N. Quincy Street Arlington, Virginia 22217			10. SOURCE OF FUNDING NUMBERS		
			PROGRAM ELEMENT NO.	PROJECT NO.	TASK NO.
			WORK UNIT ACCESSION NO.		
11. TITLE (Include Security Classification) Squeezing of Many-Atom Radiation in an Optical Cavity					
12. PERSONAL AUTHOR(S) Fu-li Li, Xiao-shen Li, D. L. Lin and Thomas F. George					
13a. TYPE OF REPORT		13b. TIME COVERED FROM _____ TO _____		14. DATE OF REPORT (Year, Month, Day) November 1989	
15. PAGE COUNT 28					
16. SUPPLEMENTARY NOTATION Prepared for publication in Physical Review A					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP	SQUEEZING SU(2) LIE ALGEBRA		
			MANY-ATOM RADIATION FOURTH ORDER		
			OPTICAL CAVITY MANY-ATOM MICROMASERS		
19. ABSTRACT (Continue on reverse if necessary and identify by block number)					
<p>A method developed in terms of the SU(2) Lie algebra for the treatment of many-level atoms is now modified to treat the system of many two-level atoms interacting with a cavity field. Special attention is given to field squeezing up to fourth order. The effects of the number of atoms and detuning on the squeezing are investigated numerically for various initial field intensities. It is found that the atomic cooperation enhances the squeezing, while in the case of weak excitation, it weakens the squeezing. The possibility of constructing many-atom micromasers is suggested.</p>					
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION Unclassified		
22a. NAME OF RESPONSIBLE INDIVIDUAL Dr. David L. Nelson			22b. TELEPHONE (Include Area Code) (202) 696-4410		22c. OFFICE SYMBOL

Squeezing of many-atom radiation in an optical cavity

Fu-li Li
Department of Physics
Xian Jiao Tong University
Xian, Shaanxi
People's Republic of China

Xiao-shen Li*, D. L. Lin and Thomas F. George
Department of Physics and Astronomy
State University of New York at Buffalo
Buffalo, New York 14260

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Special
A-1	

Abstract

A method developed in terms of the $SU(2)$ Lie algebra for the treatment of many-level atoms is now modified to treat the system of many two-level atoms interacting with a cavity field. Special attention is given to field squeezing up to fourth order. The effects of the number of atoms and detuning on the squeezing are investigated numerically for various initial field intensities. It is found that the atomic cooperation effect on the squeezing is closely related to the initial field excitation. When the initial excitation is strong, the cooperation enhances the squeezing, while in the case of weak excitation, it weakens the squeezing. The possibility of constructing many-atom micromasers is suggested.

1988 PACS Nos: 32.80.Wr, 42.50.Bs, 42.50Hz, 42.50.Md

* On leave of absence from the Shanghai Institute of Metallurgy, Chinese Academy of Sciences, Shanghai, People's Republic of China

I. Introduction

In the squeezed state, a quadrature of the electric field can have smaller quantum fluctuation than in the coherent state.¹ The potential application of such squeezed light in low-noise communications and high-precision measurements has prompted a great deal of research in recent years.

By squeezing one usually means that the mean square deviation of a certain quadrature of the field is smaller than its corresponding value when the field is in the coherent state. It has recently been generalized by Hong and Mandel,² who introduced the concept of higher-order squeezing and demonstrated its existence in such processes as degenerate parametric down conversion, resonance fluorescence and harmonic generation.³ In some cases, the higher-order squeezing may even be stronger than the ordinary second-order squeezing. In fact, the existence of fourth-order squeezing phenomena has been confirmed in theoretical studies of the single-photon and multiphoton Jaynes-Cummings (JC) model,⁴ and in the anharmonic field.⁵

Even more interesting is the fact that in the multiphoton JC model there exists intrinsic higher-order squeezing.³ Thus, higher-order squeezing has physical meaning and is not just a mathematical concept. As experimental techniques for measuring higher-order correlations in quantum optics are developing, it is significant to discuss high-order moments of the radiation field and the possibility of reduced fluctuation. It is expected from such discussions that higher-order correlation measurements may lead to the possibility of extracting the information more effectively from an optical signal.

The realization of the one-atom maser⁶ makes it possible to observe predictions of the JC-model. One of the key factors that helps keep the micromaser in continuous operation is its ability to achieve the highest

possible Rabi frequency of the atom in the optical cavity. This is because faster Rabi oscillations allow more rapid energy exchange of photons between the atom and the cavity field. The Rabi frequency can be increased by increasing the number of interacting atoms with the cavity field.⁷ In a micromaser, it is a relatively simple matter to adjust the current intensity of the atomic beam. When the beam intensity is sufficiently strong, there will be a bunch of atoms interacting simultaneously with the cavity field. It is therefore interesting to study an N -atom system interacting with a cavity field.

As the output power of a micromaser increases with increasing number of atoms, it appears that the many-atom micromaser is more attractive than the one-atom maser for practical purposes. If the mean interparticle distance is much smaller than the wave length of the cavity field, the system may be well-described by the Dicke model.⁸ A study of the many-atom micromaser on the basis of this model is being carried out and will be published in the near future. It is hoped that such studies will induce the interest of experimentalists to develop many-atom micromasers as a powerful tool for the investigation of nonclassical phenomena of the cavity field.

Radiation processes from a system of many two-level atoms in a damped cavity have been examined recently for different initial conditions.⁹ Collective effects on such properties as the atomic inversion, mean photon number and spontaneous emission caused by the presence of many atoms are explored. The purpose of this paper is to investigate how the atomic number and the strength of the coherent initial field influence fourth-order as well as second-order squeezing of the field, although second-order squeezing in the Dicke model has been briefly discussed.¹⁰

In Sec. II, we develop the theory in terms of the SU(2) Lie algebra. Numerical results are presented and discussed in Secs. III and IV. Finally, a brief discussion is given in Section V.

II. Theory

We consider a system consisting of a single-mode cavity field interacting with N atoms, each of which has two nondegenerate levels $|+\rangle$ and $|-\rangle$ with corresponding energies $\hbar\omega_+$ and $\hbar\omega_-$. Every atom is assumed to interact with the field of frequency Ω through a dipole transition only. For simplicity, we also assume that the mean interatomic distance is much smaller than the wavelength of the cavity field, so that all the atoms see exactly the same field. On the other hand, the atoms are still well separated such that their wave functions do not overlap and they see each other only through the coupling with the cavity field. Furthermore, we consider only single valence electron transitions in every atom. This means that the atoms can be treated as fermions.

The total Hamiltonian of our system can be written as

$$H = \sum_{\alpha=+,-} \hbar\omega_{\alpha} A_{\alpha}^{\dagger} A_{\alpha} + \hbar\Omega a^{\dagger} a + \hbar\lambda (A_{+}^{\dagger} A_{-} a + A_{-}^{\dagger} A_{+} a^{\dagger}) \quad (1)$$

where we have defined

$$A_{\alpha}^{\dagger} A_{\alpha} = \sum_{i=1}^N A_{\alpha}^{\dagger}(i) A_{\alpha}(i) \quad (2a)$$

$$A_+^\dagger A_- = \sum_{i=1}^N A_+^\dagger(i) A_-(i) \quad (2b)$$

$$A_-^\dagger A_+ = \sum_{i=1}^N A_-^\dagger(i) A_+(i) \quad (2c)$$

The operator A_α^\dagger (A_α) creates (annihilates) an atom in the α -level, and i labels the atom. These operators obey the anti-commutation relation

$$[A_\alpha(i), A_\beta^\dagger(j)]_+ = \delta_{ij} \delta_{\alpha\beta} \quad (3)$$

We now define, for convenience, the operators

$$N_\alpha = A_\alpha^\dagger A_\alpha \quad (4a)$$

$$D^\dagger = A_+^\dagger A_- \quad (4b)$$

$$D = A_-^\dagger A_+ \quad (4c)$$

It is a simple matter to verify that the operators defined in (4) satisfy the following commutators:

$$[N_-, D^\dagger] = -D^\dagger \quad (5a)$$

$$[N_+, D^\dagger] = D^\dagger \quad (5b)$$

$$[D, D^\dagger] = N_- - N_+ \quad (5c)$$

Equations (5) characterize the $U(2)$ Lie algebra. The first-rank Casimir operator of this algebra is just the total number operator of atoms

$$N = N_+ + N_- \quad (6)$$

It is easily seen that $[N, H] = 0$, i.e., the total number of atoms is a conserved quantity. Thus the operators N_+, N_- , D^\dagger and D defined in (4) constitute the $SU(2)$ algebra.

Since N is conserved, it may be regarded as a constant in H . Therefore, the Hamiltonian H is practically composed of the $SU(2)$ Lie algebra operators and photon operators. Consequently, if the wave function of the N -atom system belongs to a certain irreducible representation initially, it must remain in that same irreducible representation.

In the present paper, we limit our discussion to the case in which all the N atoms are initially in upper states. The atomic wave function corresponding to this case belongs to the full symmetry representation of $SU(2)$, and the state of the atomic system remains in this irreducible representation all the time. The Hilbert space of $SU(2)$ full symmetry representation can be defined by the basis vectors $|n_+, n_->$ which represent states with n_+ atoms in the upper level and n_- atoms in the lower level. Thus

$$N_+ |n_+, n_-> = n_+ |n_+, n_-> \quad (7a)$$

$$N_- |n_+, n_-> = n_- |n_+, n_-> \quad (7b)$$

$$N = n_+ + n_- \quad (7c)$$

In the space of this SU(2) full symmetry representation, the normalized state vector with an arbitrary number of atoms in the upper level can be derived, as is shown in the Appendix, from the initial state $|N,0\rangle$. The result is

$$|N-m,m\rangle = \sqrt{\frac{(N-m)!}{N!m!}} D^m |N,0\rangle, \quad 0 \leq m \leq N. \quad (8)$$

It can easily be verified that the system does not change its energy with interchange of any pair of atoms in the state.

We now introduce the total excitation operator

$$\mathcal{N} = N_+ + a^\dagger a, \quad (9)$$

which commutes with the Hamiltonian and hence represents a conserved quantity. If initially the atoms are in the state $|N,0\rangle$ and the field is in $|n\rangle$, then at any time $t > 0$ the atom-field system can only be found in a state in that part of the Hilbert space spanned by the basis vectors

$$|N-m,m\rangle |n+m\rangle. \quad (10)$$

Any arbitrary state in this subspace is characterized by the same eigenvalue $N+n$ of the operator \mathcal{N} . Hence the Hilbert space can be divided into subspaces according to the total excitation number. All the state vectors in each subspace correspond to the same eigenvalue of \mathcal{N} . Consequently, the Hamiltonian operator can never connect states from different subspaces, and we can diagonalize the Hamiltonian in each subspace.

In the subspace defined by the total excitation number $N+n$, an arbitrary state $|\phi_n\rangle$ can be expressed as

$$|\phi_n\rangle = \sum_{m=0}^N C_{nm} |N-m, m\rangle |n+m\rangle \quad (11)$$

where the expansion coefficients C_{nm} satisfy the stationary-state Schrodinger equation

$$\sum_{m'=0}^N (H_{mm'}^{(0)} + H_{mm'}^{(1)} - E\delta_{mm'}) C_{mm'} = 0 \quad (12)$$

The matrix elements in Eq. (12) are given by

$$H_{mm} = [N\hbar\omega_+ + (N+n)\hbar\Omega - (N-m)\hbar\Delta] \quad (13)$$

$$H_{mm'} = \hbar\lambda\sqrt{(n+m+1)(N-m)(m+1)} \delta_{m+1, m'} + \hbar\lambda\sqrt{(n+m'+1)(N-m')(m'+1)} \delta_{m, m'+1} \quad (14)$$

where the detuning parameter is defined by

$$\Delta = \Omega - (\omega_+ - \omega_-) \quad (15)$$

The solution of Eq. (12) provides us with energy eigenvectors and corresponding eigenvalues

$$H|\phi_{n\sigma}\rangle = E_{n\sigma}|\phi_{n\sigma}\rangle \quad (16)$$

$$|\phi_{n\sigma}\rangle = \sum_{m=0}^N c_{nm}^{\sigma} |N-m, m\rangle |n+m\rangle \quad (17)$$

where σ labels the eigenvectors in this particular subspace. The orthonormality relation is given by

$$\sum_{m=0}^N c_{nm}^{\sigma'} c_{nm}^{\sigma} = \delta_{\sigma\sigma'} \quad (18)$$

It is important to note that if initially the atomic system is in a state $|N, 0\rangle$ and the field in a linearly superimposed Fock state, then at any time $t > 0$ the atom-field system can only be found in one of the subspaces with the total excitation $N, N+1, \dots, N+n, \dots$. This can be summarized by the completeness condition

$$\sum_{n=\sigma}^{\infty} \sum_{\sigma=1}^{N+1} |\phi_{n\sigma}\rangle \langle \phi_{n\sigma}| = 1 \quad (19)$$

The equation of motion for the matrix element of the density matrix can be obtained from making use of (17) and (19), namely,

$$\frac{\partial}{\partial t} \rho_{n\sigma, n'\sigma'}(t) = -\frac{i}{\hbar} (E_{n\sigma} - E_{n'\sigma'}) \rho_{n\sigma, n'\sigma'} \quad (20)$$

The solution of (20) can be put in the form

$$\rho_{n\sigma, n'\sigma'}(t) = \rho_{n\sigma, n'\sigma'}(0) e^{-\frac{i}{\hbar} (E_{n\sigma} - E_{n'\sigma'}) t} \quad (21)$$

where the initial density matrix elements is given by

$$\rho_{n\sigma, n'\sigma'}(0) = \rho_{nn'} C_{n\sigma}^{-1} C_{n'\sigma'}^{-1} \quad (22)$$

Here, C^{-1} is the inverse matrix of the transformation defined in (11), and $\rho_{nn'}$ is the initial density matrix element of the field in the photon number representation.

Now the expectation value of any function $G(a^\dagger, a)$ can be calculated by means of the density matrix in the usual manner, namely

$$\langle G \rangle = \text{Tr}[\rho G] = \sum_n \sum_\sigma \rho_{n\sigma, n'\sigma'}(t) \langle \phi_{n'\sigma'} | G(a^\dagger, a) | \phi_{n\sigma} \rangle \quad (23)$$

To study the squeezing of the field, we introduce as usual the slowly-varying complex amplitudes of the cavity field,

$$d_1 = \frac{1}{2} [ae^{i\Omega t} + a^\dagger e^{-i\Omega t}] \quad (24a)$$

$$d_2 = \frac{1}{2i} [ae^{i\Omega t} - a^\dagger e^{-i\Omega t}] \quad (24b)$$

These operators satisfy the commutation relation

$$[d_1, d_2] = \frac{i}{2} \quad (25)$$

which implies the uncertainty relation

$$\langle (\Delta d_1)^2 \rangle \langle (\Delta d_2)^2 \rangle \geq \frac{1}{16} \quad (26)$$

The electromagnetic field is squeezed to second order if $\langle(\Delta d_1)^2\rangle < \frac{1}{4}$ or $\langle(\Delta d_2)^2\rangle < \frac{1}{4}$.

As we have mentioned above, the concept of squeezing has been generalized to K-th order where K is limited to an even number:² that is, the field is squeezed to K-th order if

$$\langle(\Delta d_{1,2})^K\rangle < (K-1)!!/2^K. \quad (27)$$

In particular, the fourth-order variance is given by

$$\langle(\Delta d_i)^4\rangle = \langle d_i^4\rangle - 4\langle d_i\rangle\langle d_i^3\rangle + 6\langle d_i^2\rangle\langle d_i\rangle^2 - 3\langle d_i\rangle^4, \quad (28)$$

and fourth-order squeezing occurs whenever $\langle(d_1)^4\rangle < \frac{3}{16}$ or $\langle(\Delta d_2)^4\rangle < \frac{3}{16}$.

III. Second-order squeezing

Throughout this paper, we take $1/\lambda$ as the unit of time and set $\hbar = 1$ in our numerical work. For the initial condition, we have assumed in the above derivation that all atoms are in their upper states. We now assume further that the field is in the coherent state at $t = 0$. Thus

$$\rho_{nn'} = \frac{z^{n'} z^{*n}}{\sqrt{n!n'!}} e^{-\bar{n}}, \quad (29)$$

where $\bar{n} = |z|^2$ is the initial mean photon number. For simplicity, we choose z to be real in this paper.

The mean square deviation of d_1 is computed as a function of time for different values of \bar{n} and Δ , the detuning parameter. For the case of

resonance interaction, we have actually calculated $\langle(\Delta d_1)^2\rangle$ for $\bar{n} = 5, 15, 25$ and 35, and the results for $\bar{n} = 5$ and 35 are plotted in Fig. 1. We observe that the squeezing becomes shallow as the number of atoms increases for $\bar{n} = 5$ or for weak excitations. When $N = 20$, the squeezing disappears completely, as is clearly seen in Fig. 1(a). The situation is very different for stronger excitations. For $\bar{n} \geq 15$, we find that the maximum squeezing increases with increasing N after the field changes for the first time from the coherent state to the squeezing state. This is illustrated, as an example, in Fig. 1(b) for $\bar{n} = 35$. Therefore, the influence of the atomic cooperative effect on second-order squeezing after the interaction is switched on is quite closely related to the initial intensity of the cavity field.

In Fig. 2, we plot the time evolution of $\langle(\Delta d_1)^2\rangle$ for various \bar{n} and $N = 10$. It is seen that during the first two squeezing periods, smaller \bar{n} corresponds to deeper squeezing of the field. However, the opposite is true during the third squeezing period, and the squeezing deepens with increasing \bar{n} , although the strongest squeezing occurs in the second period. Thus the squeezing can not be increased arbitrarily by increasing the initial field intensity, even though it is possible to cause the field to return to the squeezing state more times.

The long-time behavior of $\langle(\Delta d_1)^2\rangle$ is shown in Fig. 3 for one-atom and two-atom systems. We find that in the one-atom case the squeezing effect can be very strong at a much later time. As a matter of fact, it has been shown¹¹ that the squeezing increases with increasing \bar{n} . In the two-atom case, however, the field enters the squeezed state only briefly at a time shortly after the interaction with the atom takes place. Thus the cooperation among atoms changes the field squeezing property indeed. It is worth pointing out that although the second-order squeezing has been calculated in Ref. 8 with

the same model, no discussion of the long-time behavior or the effect due to the number of atoms was given in that work.

We have also investigated the relation between the cooperative effect and the detuning parameter. The minimum $\langle(\Delta d_1)^2\rangle$ is plotted versus Δ in Fig. 4 with fixed \bar{n} for different numbers of atoms in the cavity. It shows that for a single-atom, there exists a non-zero Δ which leads to the strongest squeezing. In cases of more than one atom in the cavity, maximum squeezing occurs only at $\Delta = 0$. This means that the cooperative interaction between the atoms weakens the contribution of the virtual atomic transition to the squeezing of the cavity field.

IV. Fourth-order squeezing

To study fourth-order squeezing, we have calculated $\langle(\Delta d_1)^4\rangle$ as a function of time for $\bar{n} = 5, 15, 25$ and 35 , and only the results for $\bar{n} = 5$ and 35 are presented here. Since there is no sign of squeezing in the long-time regime, we plot only the short-time behavior of the evolution.

Figure 5(a) shows no fourth-order squeezing at any time. Furthermore, the first minimum value of the curve increases with increasing N . This is very different from the one-atom case, in which the field can be found in the fourth-order squeezing state even if \bar{n} is as small as 0.2 .³ This indicates that the atomic cooperative effect due to the interaction with the cavity field tends to increase the fourth-order fluctuation of the field when the initial field intensity is weak. When the initial field intensity is strong, as shown in Fig. 5(b) for $\bar{n} = 35$, the first squeezing deepens with increasing N . Thus the atomic cooperative effect will reduce the fourth-order fluctuation of the field after the interaction is switched on. Therefore, the atomic cooperative effect on the field squeezing depends strongly upon the

initial field strength in the cavity. We also note that for a given \bar{n} , the field enters the squeezed state from the initial coherent state at about the same time, and this time is insensitive to the number of atoms in the cavity.

In order to study the dependence of the squeezing on the initial field excitation, we plot in Fig. 6 the time evolution of $\langle(\Delta d_1)^4\rangle$ for a given N but different \bar{n} . It is clearly seen from these curves that the first squeezing becomes shallower as \bar{n} increases, while the third squeezing deepens with increasing \bar{n} . In the case of many atoms in a cavity, it is not possible to increase the squeezing arbitrarily by increasing the initial field intensity. By so doing one can only increase the number of times the field becomes squeezed.

The influence of the variation in the detuning parameter on the fourth-order squeezing can be changed by the cooperation of atoms in the cavity. In Fig. 7, we plot the minimum of $\langle(\Delta d_1)^4\rangle$ versus the detuning for various N . Like the second-order squeezing, we find that the atomic cooperation changes the dependence of the squeezing on the detuning; that is, the squeezing reduces with increasing detuning when there is more than one atom in the cavity. On the other hand, as curve a indicates, the squeezing can be greatly increased by suitably choosing the detuning in the case of a single atom in the cavity. It is also noted that similar to the situation in Fig. 3(a) for the one-atom case, strong fourth-order squeezing occurs in the long-time regime. This phenomena disappears totally, however, for $N \geq 2$. Finally, it should be of interest to point out that fourth-order squeezing is not intrinsic to the system in the present model because it always accompanies second-order squeezing in this model.

V. Discussion

We have studied higher-order squeezing of the cavity field which is interacting with N two-level atoms by means of the $SU(2)$ Lie algebra. The method of approach is similar to what we have applied in our analysis of an M -level atom interacting with cavity fields.¹² Detailed calculations reveal that the initial field intensity plays a key role in the determination of the influence of the many-body cooperation on the field squeezing. For strong initial excitations, the atomic cooperation enhances the squeezing, while the opposite is true when the initial excitation is weak.

In a micromaser,⁶ inversely populated atoms are injected one by one into an optical cavity to create stimulated emission. We suggest that a many-atom maser may be designed in which a group of active atoms instead of one is injected into the cavity at one time. If such many-atom micromasers are realized, it should be much easier to observe the field squeezing with strong initial excitation. Furthermore, it is known that a micromaser may be expected to produce low-noise microwave radiation. If the active medium consists of only one atom, its output power is too small to be useful for any practical purpose. If, however, the active medium consists of a group of atoms, or a large number of atoms is allowed to pass the cavity at one time, the output power of the micromaser will be greatly increased. The investigation of a many-atom micromaser is being carried out and will be reported in the future.

Acknowledgments

This research was partially supported by the National Science Foundation under Grant No. CHE-8620274^{and} the Office of Naval Research.

Appendix: Derivation of Eq. (8)

To derive the state vector for an arbitrary number of atoms in the upper level, we start with the initial state

$$|N,0\rangle = \prod_{i=1}^N A_+^\dagger(i) |0\rangle, \quad (A1)$$

where $|0\rangle$ is the vacuum state defined by

$$\begin{aligned} A_+(i) |0\rangle &= 0 \\ A_-(i) |0\rangle &= 0 \end{aligned} \quad i = 1, 2, \dots, N. \quad (A2)$$

$|N,0\rangle$ is the highest-weight state from which we can obtain any arbitrary state specified by the total number of atoms N in the $SU(2)$ space of the full symmetry representation by applying the lowering operator D repeatedly. Thus we have, after m applications of the operator D , $DD\dots D|N,0\rangle = D^m|N,0\rangle$.

From the commutators

$$\begin{aligned} [N_+, D] &= -D \\ [N_-, D] &= D \end{aligned} \quad (A3)$$

we can easily find the recursion formula

$$N_+ D^m = D^m N_+ - m D^m \quad (A4)$$

$$N_- D^m = D^m N_- + m D^m$$

With the help of the definitions of number operators

$$N_+ |N,0\rangle = N |N,0\rangle \quad (A5)$$

$$N_- |N, 0\rangle = 0 \quad ,$$

it follows from Eqs. (A4) that

$$N_+ D^m |N, 0\rangle = (N-m) D^m |N, 0\rangle \quad (A6)$$

$$N_- D^m |N, 0\rangle = m D^m |N, 0\rangle \quad .$$

This means that there are $N-m$ atoms in the upper level and m atoms in the lower level in the state $D^m |N, 0\rangle$; that is,

$$D^m |N, 0\rangle \propto |N-m, m\rangle \quad , \quad 0 \leq m \leq N \quad . \quad (A7)$$

Equation (A7) represents the set of $N+1$ basis vectors that span the $SU(2)$ space of full symmetry representation corresponding to states of N atoms.

The normalization constant C can be determined in the following manner.

Since

$$D^\dagger |N, 0\rangle = A_+^\dagger A_- |N, 0\rangle = 0 \quad , \quad (A8)$$

we have

$$C^2 = \langle 0, N | (D^\dagger)^m D^m |N, 0\rangle$$

$$= \begin{cases} 1, & m = 0 \\ (N-m+1) m \langle 0, N | (D^\dagger)^{m-1} D^{m-1} |N, 0\rangle, & m \geq 1 \end{cases} \quad (A9)$$

where we have made use of the commutation relation

$$[D, D^\dagger] = N_- - N_+ \quad . \quad (A10)$$

Equation (A9) implies that

$$C^2 = N! \cdot m! / (N-m)! \quad . \quad (A11)$$

Combining (A11) with (A7), we obtain

$$|N-m, m\rangle = \sqrt{\frac{(N-m)!}{N!m!}} D^m |N, 0\rangle \quad , \quad (A12)$$

which is just Eq. (8).

References

1. H. P. Yuen, Phys. Rev. A 13, 2226 (1976).
2. C. K. Hong and L. Mandel, Phys. Rev. Lett. 54, 323 (1985); C. K. Hong and L. Mandel, Phys. Rev. A 32, 974 (1985).
3. M. Kozierowski, Phys. Rev. A 34, 3474 (1986).
4. F. L. Kien, M. Kozierowski and T. Quang, Phys. Rev. A 38, 263 (1988); L. Christopher, C. Gerry and P. J. Moyer, Phys. Rev. A 38, 5665 (1988).
5. C. Gerry and S. Rodrigues, Phys. Rev. A 35, 4440 (1987).
6. D. Meschede, H. Walther and G. Muller, Phys. Rev. Lett. 54, 551 (1985); G. Rempe, H. Walther and N. Klein, Phys. Rev. Lett. 58, 353 (1987).
7. S. M. Barnett and P. L. Knight, Optica Acta 31, 435 (1984).
8. R. H. Dicke, Phys. Rev. 93, 99 (1954).
9. J. Seke and F. Rattay, J. Opt. Soc. Am. B 4, 380 (1987); Phys. Rev. A 39, 171 (1989).
10. M. Butler and D. D. Drummond, Optica Acta 33, 1 (1986).
11. J. R. Kuklinsky and J. L. Madajczyk, Phys. Rev. A 37, 3175 (1988).
12. F. L. Li, D. L. Lin, T. F. George and X. S. Li, Phys. Rev. A 40, 1394 (1989).

Figure Captions

1. Time evolution of $\langle(\Delta d_1)^2\rangle$ for different numbers N of atoms (as indicated by the number next to each curve) and the detuning $\Delta = 0$. (a) $\bar{n} = 5$ (b) $\bar{n} = 35$.
2. Time evolution of $\langle(\Delta d_1)^2\rangle$ for various choices of the initial mean photon number \bar{n} (as indicated by the number next to each curve). Other parameters are $N = 10$ and $\Delta = 0$.
3. Long-time behavior of $\langle(\Delta d_1)^2\rangle$ for $\bar{n} = 35$ and $\Delta = 0$. (a) $N = 1$ (b) $N = 2$.
4. Minimum $\langle(\Delta d_1)^2\rangle$ versus the detuning for $\bar{n} = 25$. (a) $N = 1$, (b) $N = 2$, (c) $N = 5$.
5. Time evolution of $\langle(\Delta d_1)^4\rangle$ for different numbers N of atoms (as indicated by the number next to each curve) and $\Delta = 0$. (a) $\bar{n} = 5$, (b) $\bar{n} = 35$.
6. Time evolution of $\langle(\Delta d_1)^4\rangle$ for various choices of the initial mean photon number \bar{n} (as indicated by the number next to each curve). Other parameters are $N = 10$ and $\Delta = 0$.
7. Minimum $\langle(\Delta d_1)^4\rangle$ versus detuning for $\bar{n} = 25$. (a) $N = 1$, (b) $N = 2$, (c) $N = 5$.

Fig. 1






Fig. 2










Fig. 1









Fig. 1



TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Attn: Code 1113 800 N. Quincy Street Arlington, Virginia 22217-5000	2	Dr. David Young Code 334 NORDA NSTL, Mississippi 39529	1
Dr. Bernard Douda Naval Weapons Support Center Code 50C Crane, Indiana 47522-5050	1	Naval Weapons Center Attn: Dr. Ron Atkins Chemistry Division China Lake, California 93555	1
Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko, Code L52 Port Hueneme, California 93401	1	Scientific Advisor Commandant of the Marine Corps Code RD-1 Washington, D.C. 20380	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12 high quality	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, NC 27709	1
DTNSRDC Attn: Dr. H. Singerman Applied Chemistry Division Annapolis, Maryland 21401	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1
Dr. William Tolles Superintendent Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375-5000	1	Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 91232	1
		Dr. David L. Nelson Chemistry Division Office of Naval Research 800 North Quincy Street Arlington, Virginia 22217	1

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. J. E. Jensen
Hughes Research Laboratory
3011 Malibu Canyon Road
Malibu, California 90265

Dr. C. B. Harris
Department of Chemistry
University of California
Berkeley, California 94720

Dr. J. H. Weaver
Department of Chemical Engineering
and Materials Science
University of Minnesota
Minneapolis, Minnesota 55455

Dr. F. Kutzler
Department of Chemistry
Box 5055
Tennessee Technological University
Cookeville, Tennessee 38501

Dr. A. Reisman
Microelectronics Center of North Carolina
Research Triangle Park, North Carolina
27709

Dr. D. DiLella
Chemistry Department
George Washington University
Washington D.C. 20052

Dr. M. Grunze
Laboratory for Surface Science and
Technology
University of Maine
Orono, Maine 04469

Dr. R. Reeves
Chemistry Department
Rensselaer Polytechnic Institute
Troy, New York 12181

Dr. J. Butler
Naval Research Laboratory
Code 6115
Washington D.C. 20375-5000

Dr. Steven M. George
Stanford University
Department of Chemistry
Stanford, CA 94305

Dr. L. Interante
Chemistry Department
Rensselaer Polytechnic Institute
Troy, New York 12181

Dr. Mark Johnson
Yale University
Department of Chemistry
New Haven, CT 06511-8118

Dr. Irvin Heard
Chemistry and Physics Department
Lincoln University
Lincoln University, Pennsylvania 19352

Dr. W. Knauer
Hughes Research Laboratory
3011 Malibu Canyon Road
Malibu, California 90265

Dr. K.J. Klaubunde
Department of Chemistry
Kansas State University
Manhattan, Kansas 66506

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. G. A. Somorjai
Department of Chemistry
University of California
Berkeley, California 94720

Dr. J. Murday
Naval Research Laboratory
Code 6170
Washington, D.C. 20375-5000

Dr. J. B. Hudson
Materials Division
Rensselaer Polytechnic Institute
Troy, New York 12181

Dr. Theodore E. Madey
Surface Chemistry Section
Department of Commerce
National Bureau of Standards
Washington, D.C. 20234

Dr. J. E. Demuth
IBM Corporation
Thomas J. Watson Research Center
P.O. Box 218
Yorktown Heights, New York 10598

Dr. M. G. Lagally
Department of Metallurgical
and Mining Engineering
University of Wisconsin
Madison, Wisconsin 53706

Dr. R. P. Van Duyne
Chemistry Department
Northwestern University
Evanston, Illinois 60637

Dr. J. M. White
Department of Chemistry
University of Texas
Austin, Texas 78712

Dr. D. E. Harrison
Department of Physics
Naval Postgraduate School
Monterey, California 93940

Dr. R. L. Park
Director, Center of Materials
Research
University of Maryland
College Park, Maryland 20742

Dr. W. T. Peria
Electrical Engineering Department
University of Minnesota
Minneapolis, Minnesota 55455

Dr. Keith H. Johnson
Department of Metallurgy and
Materials Science
Massachusetts Institute of Technology
Cambridge, Massachusetts 02139

Dr. S. Sibener
Department of Chemistry
James Franck Institute
5640 Ellis Avenue
Chicago, Illinois 60637

Dr. Arnold Green
Quantum Surface Dynamics Branch
Code 3817
Naval Weapons Center
China Lake, California 93555

Dr. A. Wold
Department of Chemistry
Brown University
Providence, Rhode Island 02912

Dr. S. L. Bernasek
Department of Chemistry
Princeton University
Princeton, New Jersey 08544

Dr. W. Kohn
Department of Physics
University of California, San Diego
La Jolla, California 92037

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. F. Carter
Code 6170
Naval Research Laboratory
Washington, D.C. 20375-5000

Dr. Richard Colton
Code 6170
Naval Research Laboratory
Washington, D.C. 20375-5000

Dr. Dan Pierce
National Bureau of Standards
Optical Physics Division
Washington, D.C. 20234

Dr. R. Stanley Williams
Department of Chemistry
University of California
Los Angeles, California 90024

Dr. R. P. Messmer
Materials Characterization Lab.
General Electric Company
Schenectady, New York 22217

Dr. Robert Gomer
Department of Chemistry
James Franck Institute
5640 Ellis Avenue
Chicago, Illinois 60637

Dr. Ronald Lee
R301
Naval Surface Weapons Center
White Oak
Silver Spring, Maryland 20910

Dr. Paul Schoen
Code 6190
Naval Research Laboratory
Washington, D.C. 20375-5000

Dr. John T. Yates
Department of Chemistry
University of Pittsburgh
Pittsburgh, Pennsylvania 15260

Dr. Richard Greene
Code 5230
Naval Research Laboratory
Washington, D.C. 20375-5000

Dr. L. Kesmodel
Department of Physics
Indiana University
Bloomington, Indiana 47403

Dr. K. C. Janda
University of Pittsburgh
Chemistry Building
Pittsburg, PA 15260

Dr. E. A. Irene
Department of Chemistry
University of North Carolina
Chapel Hill, North Carolina 27514

Dr. Adam Heller
Bell Laboratories
Murray Hill, New Jersey 07974

Dr. Martin Fleischmann
Department of Chemistry
University of Southampton
Southampton SO9 5NH
UNITED KINGDOM

Dr. H. Tachikawa
Chemistry Department
Jackson State University
Jackson, Mississippi 39217

Dr. John W. Wilkins
Cornell University
Laboratory of Atomic and
Solid State Physics
Ithaca, New York 14853

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. R. G. Wallis
Department of Physics
University of California
Irvine, California 92664

Dr. D. Ramaker
Chemistry Department
George Washington University
Washington, D.C. 20052

Dr. J. C. Hemminger
Chemistry Department
University of California
Irvine, California 92717

Dr. T. F. George
Chemistry Department
University of Rochester
Rochester, New York 14627

Dr. G. Rubloff
IBM
Thomas J. Watson Research Center
P.O. Box 218
Yorktown Heights, New York 10598

Dr. Horia Metiu
Chemistry Department
University of California
Santa Barbara, California 93106

Dr. W. Goddard
Department of Chemistry and Chemical
Engineering
California Institute of Technology
Pasadena, California 91125

Dr. P. Hansma
Department of Physics
University of California
Santa Barbara, California 93106

Dr. J. Baldeschwieler
Department of Chemistry and
Chemical Engineering
California Institute of Technology
Pasadena, California 91125

Dr. J. T. Keiser
Department of Chemistry
University of Richmond
Richmond, Virginia 23173

Dr. R. W. Plummer
Department of Physics
University of Pennsylvania
Philadelphia, Pennsylvania 19104

Dr. E. Yeager
Department of Chemistry
Case Western Reserve University
Cleveland, Ohio 41106

Dr. N. Winograd
Department of Chemistry
Pennsylvania State University
University Park, Pennsylvania 16802

Dr. Roald Hoffmann
Department of Chemistry
Cornell University
Ithaca, New York 14853

Dr. A. Stecki
Department of Electrical and
Systems Engineering
Rensselaer Polytechnic Institute
Troy, New York 12181

Dr. G.H. Morrison
Department of Chemistry
Cornell University
Ithaca, New York 14853